



Payame Noor University



Control and Optimization in Applied Mathematics (COAM)

DOI. 10.30473/COAM.2021.59584.1165

Vol. 5, No. 2, Summer - Autumn 2020 (83-91), ©2016 Payame Noor University, Iran

Research Article

Topological Subdifferential and its Role in Nonsmooth Optimization with Quasiconvex Data

H. Soroush*

Department of Mathematics, Payame Noor University,
P.O. Box. 19395-3697, Tehran, Iran.

Received: June 17, 2021; **Accepted:** October 20, 2021.

Abstract. In this paper, we study nonsmooth optimization problems with quasiconvex functions using topological subdifferential. We present some necessary and sufficient optimality conditions and characterize topological pseudoconvex functions. Finally, the Mond-Weir type weak and strong duality results are stated for the problems.

Keywords. Quasiconvex optimization, Duality theorems, Optimality conditions, Pseudoconvex, Subdifferential.

MSC. 90C34; 90C40.

* Corresponding author
h_soroush2011@pnu.ac.ir
<http://mathco.journals.pnu.ac.ir>

1 Introduction

A mathematical optimization problem with a quasiconvex objective function and finitely many quasiconvex constraints is called a *quasiconvex programming problem*. Problems of this type have been utilized for the modeling of topological optimization problems and some theoretical topics. (see, e.g., [3, 5, 6, 7] and therein references).

This paper deals with a nonsmooth quasiconvex programming problem. We do not assume that the data of the problem are differentiable. Thus, we replace the derivative appearing in the classical results with topological subdifferential. We refer to [10] and [11, Section 5] to see the importance of this subdifferential and its relation to other subdifferentials.

It is worth to mention that some necessary conditions of Karush–Kuhn–Tucker (KKT) type for optimality of nonsmooth quasiconvex programming problems have been presented in [7] under various subdifferentials; for instance, Greenberg–Pierskalla subdifferential, Penot subdifferential, Plastria subdifferential, and Gutiérrez subdifferential. Since there are no articles that study necessary and sufficient optimality conditions under topological subdifferential, the aim of this paper is to fill this gap as the first task.

The structure of subsequent sections of this paper is as follows: In Section 2, we establish required definitions and preliminary results which are required thereafter. In Section 3, we present the KKT type necessary and sufficient optimality conditions for nonsmooth quasiconvex problems, and as applications of proved optimality conditions, we study the duality results of the problem.

2 Notations and Preliminaries

In this section, we present some definitions and auxiliary results that will be needed in what follows.

For a non-empty subset M of \mathbb{R}^n , its polar cone is defined as

$$M^0 := \{x \in \mathbb{R}^n \mid \langle x, d \rangle \leq 0, \quad \forall d \in M\},$$

where $\langle \cdot, \cdot \rangle$ denotes the standard inner-product in \mathbb{R}^n . Also, \overline{M} , $\text{conv}(M)$, and $\text{cone}(M)$ denote the closure of M , the convex hull of M , and the convex cone hull of M , respectively. Moreover, the attainable cone and the interior cone of M at $x_0 \in M$ are respectively defined as [2]:

$$\mathcal{A}(M, x_0) := \left\{ z \in \mathbb{R}^n \mid \forall t_k \downarrow 0, \exists z_k \rightarrow z, x_0 + t_k z_k \in M \quad \forall t \in \mathbb{N} \right\},$$

$$\mathcal{I}(M, x_0) := \left\{ z \in \mathbb{R}^n \mid \exists K > 0, \forall t_k \downarrow 0, \forall z_k \rightarrow z, x_0 + t_k z_k \in M, \quad \forall k \geq K \right\}.$$

Observe that $\mathcal{A}(M, x_0)$ is always a non-empty closed cone ([2, Theorem 3.4.3]) and $\mathcal{I}(M, x_0)$ is always a open cone ([2, Theorem 3.4.6]).

Theorem 1. [2, Theorem 3.4.10] Let $C \subseteq \mathbb{R}^n$ be a convex set. If $\mathcal{I}(M, x_0) \neq \emptyset$, then

$$\overline{\mathcal{I}(M, x_0)} = \mathcal{A}(M, x_0).$$

Let C be a convex subset of \mathbb{R}^n and $x_0 \in C$. The normal cone of C at x_0 is denoted by $N(C, x_0)$, i.e.,

$$N(C, x_0) := \left\{ y \in \mathbb{R}^n \mid \langle y, x - x_0 \rangle \leq 0 \quad \forall x \in C \right\}.$$

The zero vector of \mathbb{R}^n is denoted by 0_n . If $K \subseteq \mathbb{R}^n$ is a convex cone, [2, Theorem 2.3.3] and the fact that $N(K^0, 0_n) = (K^0)^0$ imply that

$$N(K^0, 0_n) = \overline{K}. \tag{1}$$

We recall from [11] that the topological (or incident, or upper epi-) directional derivative of a given function $\varphi : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ at $\hat{x} \in \text{dom}\varphi := \{x \in \mathbb{R}^n \mid \varphi(x) \in \mathbb{R}\}$ is defined by

$$\varphi^b(\hat{x}; v) := \sup_{\varepsilon > 0} \limsup_{t \downarrow 0} \inf_{\|u-v\| < \varepsilon} \frac{\varphi(\hat{x} + tu) - \varphi(\hat{x})}{t}, \quad \forall v \in \mathbb{R}^n.$$

Also, the epigraph of φ is denoted by E_φ ,

$$E_\varphi := \{(x, r) \in \mathbb{R}^n \times \mathbb{R} \mid \varphi(x) \leq r\}.$$

The following theorem presents an important property of topological directional derivative.

Theorem 2. [11, Page 263] & [2, Page 245] The epigraph of function $v \rightarrow \varphi^b(\hat{x}; v)$ coincides to attainable cone of E_φ at $(\hat{x}, \varphi(\hat{x}))$. In other word,

$$E_{\varphi^b(\hat{x}; \cdot)} = \mathcal{A}(E_\varphi, \hat{x}_\varphi),$$

in which, $\hat{x}_\varphi := (\hat{x}, \varphi(\hat{x}))$.

We say that φ is regular at \hat{x} if the function $v \rightarrow \varphi^b(\hat{x}; v)$ is convex (this type of functions, without special naming, have been studied in [9, Proposition 6.5]). According to Theorem 2 and [2, Theorem 2.5.1], we conclude that φ is regular at \hat{x} iff $\mathcal{A}(E_\varphi, \hat{x}_\varphi)$ is a convex set. Also, if φ is convex and $\hat{x} \in \text{dom}\varphi$, it is regular at \hat{x} by [12, Theorem 23.1] (see also [2, Theorem 2.5.1]). Let $r \in \mathbb{R} \cup \{+\infty\}$ be given. The r -sublevel set of $\varphi : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ is denoted by $[\varphi \leq r]$,

$$[\varphi \leq r] := \{x \in \mathbb{R}^n \mid \varphi(x) \leq r\}.$$

A function $\varphi : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ is said to be quasiconvex, if for each $r \in \mathbb{R} \cup \{+\infty\}$ its corresponding r -sublevel set $[\varphi \leq r]$ is convex. We can see that φ is quasiconvex if and only if ([2, Theorem 2.10.1])

$$\varphi(\mu x + (1 - \mu)y) \leq \max\{\varphi(x), \varphi(y)\}, \quad \forall x, y \in \mathbb{R}^n, \quad \forall \mu \in [0, 1]. \tag{2}$$

Equivalently, φ is quasiconvex if and only if

$$\left(\varphi(y) \leq \varphi(x), \quad 0 \leq \mu \leq 1 \right) \implies \varphi(\mu x + (1 - \mu)y) \leq \varphi(x).$$

The topological (or incident [11]) subdifferential of a $\varphi : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ at $\hat{x} \in \text{dom}\varphi$ is defined by

$$\partial^b \varphi(\hat{x}) := \{\xi \in \mathbb{R}^n \mid \langle \xi, v \rangle \leq \varphi^b(\hat{x}; v), \quad \forall v \in \mathbb{R}^n\}.$$

In [10, Propositions 1, 3, 16] some extraordinary properties for $\varphi^b(\hat{x}; v)$ and $\partial^b \varphi(\hat{x})$ which shows their important rule and uniqueness, were proved. For instance, see the following results:

Theorem 3. [10, Proposition 1] $\varphi^b(\hat{x}; \cdot)$ is a lower semi-continuous (l.s.c. in brief) quasiconvex function when φ is a quasiconvex function.

We are able to see ([10, Proposition 47, and Corollary 50]) that for a convex function $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$, $\partial^b \phi(\hat{x})$ coincides with the Fenchel subdifferential $\partial \phi(\hat{x})$, is defined as follows:

$$\partial \phi(\hat{x}) := \{\xi \in \mathbb{R}^n \mid \phi(x) \geq \phi(\hat{x}) + \langle \xi, x - \hat{x} \rangle, \quad \forall x \in \mathbb{R}^n\}.$$

It is worth to observe that $\partial \phi(\hat{x})$ is a non-empty closed convex set in \mathbb{R}^n ([2, Page 373]), and if \hat{x} is a minimizer of convex function ϕ on a convex set C , then ([2, Theorem 4.3.2])

$$0_n \in \partial \phi(\hat{x}) + N(C, \hat{x}). \tag{3}$$

3 Main Results

Throughout this paper, we shall consider the following optimization problem:

$$(P): \quad \inf \vartheta(x), \quad \text{s.t.} \quad x \in \Omega := \left\{x \in \mathbb{R}^n \mid \psi_j(x) \leq 0 \quad \forall j \in J := \{1, \dots, m\}\right\},$$

where ϑ and ψ_j , $j \in J$ are quasiconvex functions from \mathbb{R}^n to $\mathbb{R} \cup \{+\infty\}$.

Remark 1. The quasiconvexity of ψ_j functions (for $j \in J$) and the fact that

$$\Omega = \bigcap_{j=1}^m \{x \in \mathbb{R}^n \mid \psi_j(x) \leq 0\} = \bigcap_{j=1}^m [\psi_j \leq 0],$$

imply that the feasible set Ω is convex. Thus, $\mathcal{A}(\Omega, \hat{x}) = \overline{\mathcal{I}(\Omega, \hat{x})}$ whenever $\mathcal{I}(\Omega, \hat{x}) \neq \emptyset$ by Theorem 1.

Lemma 1. Suppose that $\hat{x} \in \Omega$ is an optimal solution of (P). If $\mathcal{I}(\Omega, \hat{x}) \neq \emptyset$ and ϑ is regular at \hat{x} , then

$$\vartheta^b(\hat{x}; v) \geq 0, \quad \forall v \in \mathcal{A}(\Omega, \hat{x}).$$

Proof. Let $d \in \mathcal{I}(\Omega, \hat{x})$. By definition of interior cone, for each sequence $d_k \rightarrow d$ and $t_k \rightarrow 0^+$, $\hat{x} + t_k d_k \in \Omega$ for all $k \in \mathbb{N}$. Thus, the optimality of \hat{x} implies that $\vartheta(\hat{x} + t_k d_k) - \vartheta(\hat{x}) \geq 0$, and hence

$$\vartheta^b(\hat{x}; d) \geq 0 \quad \forall d \in \mathcal{I}(\Omega, \hat{x}). \tag{4}$$

Considering Remark 1, for each $v \in \mathcal{A}(\Omega, \hat{x}) = \overline{\mathcal{I}(\Omega, \hat{x})}$, we can find a sequence $\{d_k\}_{k=1}^\infty$ in $\mathcal{I}(\Omega, \hat{x})$ converging to v . Taking into consideration the continuity of $\vartheta^b(\hat{x}; \cdot)$ (by the regularity of ϑ) and the validity of (4), we obtain that

$$\vartheta^b(\hat{x}; v) = \lim_{k \rightarrow \infty} \vartheta^b(\hat{x}; d_k) \geq 0,$$

and the proof is complete. □

For a given $x_0 \in \Omega$, with the convention $\bigcup_{j \in \emptyset} X_j = \emptyset$, set

$$J(x_0) := \{j \in J \mid \psi_j(x_0) = 0\},$$

$$\mathcal{Z}(x_0) := \bigcup_{j \in J(x_0)} \partial^b \psi_j(x_0).$$

The following constraint qualification will be needed for stating the necessary optimality condition in Karush-Kuhn-Tucker (KKT) type.

Definition 1. Let $x_0 \in \Omega$. We say that the generalized Kuhn-Tucker constraint qualification (GKTCQ in brief) holds at x_0 if

$$\left(\mathcal{Z}(x_0)\right)^0 \subseteq \mathcal{A}(\Omega, x_0).$$

Theorem 4. Suppose that $\hat{x} \in \Omega$ is an optimal solution of (P), $\mathcal{I}(\Omega, \hat{x}) \neq \emptyset$, and that ϑ is regular at \hat{x} . If the (GKTCQ) is satisfied at \hat{x} , then

$$0_n \in \partial^b \vartheta(\hat{x}) + \overline{\text{conv}(\mathcal{Z}(\hat{x}))}.$$

Proof. At the first, we observe from [11, Page 263] that $\vartheta^b(\hat{x}; 0_n)$ is either 0 or $-\infty$. On the other hand, because of $0 \in \mathcal{A}(\Omega, \hat{x})$, we have $\vartheta^b(\hat{x}; 0_n) \geq 0$ by Lemma 1. Thus,

$$\vartheta^b(\hat{x}; 0_n) = 0. \quad (5)$$

Let $v \in \left(\overline{\text{conv}(\mathcal{Z}(\hat{x}))}\right)^0$. Because of $\left(\overline{\text{conv}(\mathcal{Z}(\hat{x}))}\right)^0 = \left(\mathcal{Z}(\hat{x})\right)^0$ and the (GKTCQ), we conclude $\vartheta^b(\hat{x}; v) \geq 0$ in view of Lemma 1. By (5) we thus obtain that the following optimization problem has a local solution at $\bar{v} := 0_n$:

$$\begin{aligned} \min \quad & \vartheta^b(\hat{x}; \cdot)(v) \\ \text{s.t.} \quad & v \in \left(\overline{\text{conv}(\mathcal{Z}(\hat{x}))}\right)^0. \end{aligned}$$

Since the objective function and the constraint set of above problem are convex, by (3) we give

$$0_n \in \partial(\vartheta^b(\hat{x}; \cdot))(0_n) + N\left(\left(\overline{\text{conv}(\mathcal{Z}(\hat{x}))}\right)^0, 0_n\right). \quad (6)$$

Now, we observe that $N\left(\left(\overline{\text{conv}(\mathcal{Z}(\hat{x}))}\right)^0, 0_n\right) = \overline{\text{conv}(\mathcal{Z}(\hat{x}))}$ by (1), and we have also

$$\begin{aligned} \partial(\vartheta^b(\hat{x}; \cdot))(0_n) &= \left\{ \xi \in \mathbb{R}^n \mid \vartheta^b(\hat{x}; w) - \vartheta^b(\hat{x}; 0_n) \geq \langle \xi, w - 0_n \rangle \quad \forall w \in \mathbb{R}^n \right\} \\ &= \left\{ \xi \in \mathbb{R}^n \mid \vartheta^b(\hat{x}; w) \geq \langle \xi, w \rangle \quad \forall w \in \mathbb{R}^n \right\} \\ &= \partial^b \vartheta(\hat{x}). \end{aligned} \quad (7)$$

Hence, (6) implies that

$$0_n \in \partial^b \vartheta(\hat{x}) + \overline{\text{conv}(\mathcal{Z}(\hat{x}))},$$

and the proof is complete. \square

We should mention that the following KKT type necessary condition is similar that one proved in [1] and [4] using Clarke subdifferential.

Theorem 5. (KKT necessary condition) Let $\hat{x} \in \Omega$ be an optimal solution of (P). Assume that ϑ is regular at \hat{x} , $\mathcal{I}(\Omega, \hat{x}) \neq \emptyset$, and the (GKTCQ) is satisfied at \hat{x} . Moreover, if $\text{conv}(\mathcal{Z}(\hat{x}))$ is a closed set, then there exist $\lambda_j \geq 0$, $j \in J(\hat{x})$, such that

$$0_n \in \partial^b \vartheta(\hat{x}) + \sum_{j \in J(\hat{x})} \lambda_j \partial^b \psi_j(\hat{x}), \quad \text{and} \quad \sum_{j \in J(\hat{x})} \lambda_j = 1.$$

Proof. It follows from Theorem 4 and the closedness of $\text{conv}(\mathcal{Z}(\hat{x}))$ that

$$0_n \in \partial^b \vartheta(\hat{x}) + \overline{\text{conv}(\mathcal{Z}(\hat{x}))} = \partial^b \vartheta(\hat{x}) + \text{conv}(\mathcal{Z}(\hat{x})).$$

Thus, there exist $\xi^\vartheta \in \partial^b \vartheta(\hat{x})$ and $w \in \text{conv}(\mathcal{Z}(\hat{x}))$ such that

$$0_n = \xi^\vartheta + w. \quad (8)$$

On the other hand, the regularity assumption of ψ_j as $j \in J(\hat{x})$ and (7) imply that $\partial^b \psi_j(\hat{x})$ is a closed convex set in \mathbb{R}^n for $j \in J(\hat{x})$. Thus, the well-known equality (see, e.g., [12, Theorem 3.3])

$$\text{conv}(\mathcal{Z}(\hat{x})) = \left\{ \sum_{j \in J(\hat{x})} \lambda_j \zeta_j \mid \sum_{j \in J(\hat{x})} \lambda_j = 1, \lambda_j \geq 0, \zeta_j \in \partial^b \psi_j(\hat{x}), j \in J(\hat{x}) \right\},$$

implies that for each $j \in J(\hat{x})$ there exist $\zeta_j \in \partial^b \psi_j(\hat{x})$ and $\lambda_j \geq 0$ such that

$$w = \sum_{j \in J(\hat{x})} \lambda_j \zeta_j, \quad \text{and} \quad \sum_{j \in J(\hat{x})} \lambda_j = 1.$$

The last relations and (8) allow us to conclude that

$$0_n = \xi^\vartheta + \sum_{j \in J(\hat{x})} \lambda_j \zeta_j \in \partial^b \vartheta(\hat{x}) + \sum_{j \in J(\hat{x})} \lambda_j \partial^b \psi_j(\hat{x}),$$

as required. \square

The following useful lemma characterizes quasiconvex functions using topological subdifferential.

Lemma 2. If $\varphi : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ is a quasiconvex function, then

$$\varphi(x) \leq \varphi(x_0) \implies \langle \xi, x - x_0 \rangle \leq 0 \quad \forall \xi \in \partial^b \varphi(x_0).$$

Proof. Assume that $\varphi(x) \leq \varphi(x_0)$. By (2), for each $t \in (0, 1)$, we have

$$\varphi(x_0 + t(x - x_0)) = \varphi(tx + (1-t)x_0) \leq \max\{\varphi(x), \varphi(x_0)\} = \varphi(x_0).$$

This implies $\varphi^b(x_0; x - x_0) \leq 0$. Thus, by the definition of incident subdifferential, we get

$$0 \geq \varphi^b(x_0; x - x_0) \geq \sup\{\langle \xi, x - x_0 \rangle \mid \xi \in \partial^b \varphi(x_0)\}.$$

Therefore, $\langle \xi, x - x_0 \rangle \leq 0$ for all $\xi \in \partial^b \varphi(x_0)$, as required. \square

Definition 2. The function $\varphi : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ is said to be topological pseudoconvex at x_0 if

$$\varphi(x) < \varphi(x_0) \implies \langle \xi, x - x_0 \rangle < 0 \quad \forall \xi \in \partial^b \varphi(x_0).$$

A sufficient condition for the topological pseudoconvexity of quasiconvex functions is presented in following theorem.

Theorem 6. Let $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}$ be an upper-semicontinuous quasiconvex function with $0_n \notin \partial^b \varphi(x_0)$ for some $x_0 \in \mathbb{R}^n$. Then φ is topological pseudoconvex at x_0 .

Proof. Assume that

$$\varphi(x) < \varphi(x_0). \tag{9}$$

Let ξ be an arbitrary element in $\partial^b \varphi(x_0)$. We claim that

$$\langle \xi, x - x_0 \rangle < 0. \tag{10}$$

Suppose on the contrary that $\langle \xi, x - x_0 \rangle \geq 0$. Thus, Lemma 2 implies that $\langle \xi, x - x_0 \rangle = 0$. This means that there exists a sequence $\{u_l\}_{l=1}^\infty$ such that

$$\lim_{l \rightarrow \infty} u_l = x - x_0, \quad \text{and} \quad \langle \xi, u_l \rangle > 0, \quad \forall l \in \mathbb{N}.$$

The latter can be rewritten as

$$\langle \xi, (u_l + x_0) - x_0 \rangle > 0, \quad \forall l \in \mathbb{N}.$$

From this and Lemma 2, we conclude that $\varphi(u_l + x_0) \geq \varphi(x_0)$. After passing to the limit as $l \rightarrow \infty$, we deduce that $\varphi(x) \geq \varphi(x_0)$ by the upper-semicontinuity of φ . This contradicts (9), and so, (10) holds. \square

Now, we can state a sufficient optimality condition of KKT type.

Theorem 7. (KKT sufficient condition) Let $\hat{x} \in \Omega$. Assume that ϑ is topological pseudoconvex at \hat{x} , and there exist $\lambda_j \geq 0$ for $j \in J(\hat{x})$, such that

$$0_n \in \partial^b \vartheta(\hat{x}) + \sum_{j \in J(\hat{x})} \lambda_j \partial^b \psi_j(\hat{x}). \quad (11)$$

Then \hat{x} is a global solution of (P).

Proof. Set $J(\hat{x}) = \{j_1, \dots, j_p\}$. According to (11), we can find some $\xi \in \partial^b \vartheta(\hat{x})$ and $\xi_{j_k} \in \partial^b \psi_{j_k}(\hat{x})$, $k = 1, 2, \dots, p$, such that

$$\xi + \lambda_{j_1} \xi_{j_1} + \dots + \lambda_{j_p} \xi_{j_p} = 0_n. \quad (12)$$

Suppose on the contrary that there exists some $x^* \in \Omega$ such that $\vartheta(x^*) < \vartheta(\hat{x})$. From $\xi \in \partial^b \vartheta(\hat{x})$ and the topological pseudoconvexity of ϑ at \hat{x} , we get

$$\langle \xi, x^* - \hat{x} \rangle < 0. \quad (13)$$

Since $\psi_{j_k}(x^*) \leq 0 = \psi_{j_k}(\hat{x})$ for $k = 1, 2, \dots, p$, Lemma 2 allows us to conclude that

$$\langle \lambda_{j_k} \xi_{j_k}, x^* - \hat{x} \rangle \leq 0 \quad \forall k = 1, 2, \dots, p. \quad (14)$$

Owing to the inequalities (13) and (14), we deduce that

$$\langle \xi + \lambda_{j_1} \xi_{j_1} + \dots + \lambda_{j_p} \xi_{j_p}, x^* - \hat{x} \rangle < 0,$$

which contradicts (12). The proof is complete. \square

Now, we apply the optimality conditions which were presented in the previous theorems to give the weak and strong duality results for (P). We consider the following Mond-Weir [8] dual problem to (P):

$$(MD): \quad \max \vartheta(y) \quad \text{s.t.} \quad y \in \Lambda,$$

where Λ is defined as:

$$\Lambda := \left\{ y \in \mathbb{R}^n \mid \exists \lambda_j \geq 0 \text{ for } j \in J(y), \text{ such that } 0_n \in \partial^b \vartheta(y) + \sum_{j \in J(y)} \partial^b \lambda_j \psi_j(y) \right\}.$$

Theorem 8. (*weak duality*) Suppose that $\hat{x} \in \Omega$ and $\hat{y} \in \Lambda$. If ϑ is topological pseudoconvex at \hat{y} , then $\vartheta(\hat{x}) \geq \vartheta(\hat{y})$.

Proof. Definition of Λ and the assumption $\hat{y} \in \Lambda$ imply that there exist some $\zeta \in \partial^b \vartheta(\hat{y})$, $\lambda_j \geq 0$ and $\zeta_j \in \partial^b \psi_j(\hat{y})$ for $j \in J(\hat{y})$, such that

$$\zeta + \sum_{j \in J(\hat{y})} \lambda_j \zeta_j = 0_n. \tag{15}$$

Assume on the contrary that $\vartheta(\hat{x}) < \vartheta(\hat{y})$. So, the topological pseudoconvexity of ϑ at \hat{x} implies that

$$\langle \zeta, \hat{x} - \hat{y} \rangle < 0. \tag{16}$$

On the other hand, the quasiconvexity of ψ_j s at \hat{y} , Lemma 2, and the inequality $\psi_j(\hat{x}) \leq 0 = \psi_j(\hat{y})$ for $j \in J(\hat{y})$ imply that

$$\langle \zeta_j, \hat{x} - \hat{y} \rangle \leq 0, \quad \text{for all } j \in J(\hat{y}).$$

The above inequality, (16), and $\lambda_j \geq 0$ for $j \in J(\hat{y})$ conclude that

$$\langle \zeta + \sum_{j \in J(\hat{y})} \lambda_j \zeta_j, \hat{x} - \hat{y} \rangle = \langle \zeta, \hat{x} - \hat{y} \rangle + \sum_{j \in J(\hat{y})} \lambda_j \langle \zeta_j, \hat{x} - \hat{y} \rangle < 0,$$

which contradicts (15). The proof is complete. □

Theorem 9. (*strong duality*) Let \hat{x} and \hat{y} be respectively the optimal solutions of (P) and (MD). Assume that ϑ and ψ_j are regular at \hat{x} as $j \in J(\hat{x})$, $\mathcal{I}(\Omega, \hat{x}) \neq \emptyset$, and the (GKTCQ) is satisfied at \hat{x} . If ϑ is topological pseudoconvex at \hat{y} and $\text{conv}(\mathcal{Z}(\hat{x}))$ is a closed set, the optimal values of (P) and (MD) are equal, i.e.,

$$\vartheta(\hat{x}) = \vartheta(\hat{y}).$$

Proof. Since the hypothesis of Theorem 5 are all true, we can find some scalars $\lambda_j \geq 0$, $j \in J(\hat{x})$ such that

$$0_n \in \partial^b \vartheta(\hat{x}) + \sum_{j \in J(\hat{x})} \partial^b \lambda_j \psi_j(\hat{x}).$$

This implies that $\hat{x} \in \Lambda$, i.e., \hat{x} is a feasible point for (MD). Because of \hat{y} is optimal solution of (MD), the inequality $\vartheta(\hat{x}) \leq \vartheta(\hat{y})$ holds. This inequality and weak duality Theorem 9 imply $\vartheta(\hat{x}) = \vartheta(\hat{y})$, as required. □

References

- [1] Caristi G., Kanzi N. 2015. “Karush-Kuhn-Tucker type conditions for optimality of non-smooth multiobjective semi-infinite programming”, *International Journal of Mathematical Analysis* 9: 1929-1938.
- [2] Giorgi G., Guerraggio A., Thierselder J. 2004. “Mathematics of Optimization; Smooth and Nonsmooth cases”, *Elsivier*.
- [3] Hassani Bafrani A., Sadeghieh A. 2018. “Quasi-gap and gap functions for non-smooth multi-objective semi-infinite optimization problems”, *Control and Optimization in Applied Mathematics*. 3, 1-12.
- [4] Kanzi N. 2018. “Necessary and sufficient conditions for (weakly) efficient of nondifferentiable multi-objective semi-infinite programming”, *Iranian Journal of Science and Technology*, 3, 1537-1544.

-
- [5] Kanzi N., Sadeghieh A., Caristi G. 2019. "Optimality conditions for semi-infinite programming problems involving generalized convexity", *Optimization Letters*, 13, 113-126.
 - [6] Kanzi N., Soleymani-damaneh M. 2015. "Slater CQ, optimality and duality for quasiconvex semi-infinite optimization problems", *Journal of Mathematical Analysis and Applications*, 434: 638-651.
 - [7] Kanzi N., Soleymani-damaneh M. 2020. "Characterization of the weakly efficient solutions in nonsmooth quasiconvex multiobjective optimization", *Journal of Global Optimization*, 77: 627-641.
 - [8] Mond B., Weir T. 1981. "Generalized concavity and duality", In, Schaible. S., Ziemba, W.T. (eds). *Generalized concavity in Optimization and Economics*, 263-279.
 - [9] Penot J. P. 2000. "What is quasiconvex analysis?", *Optimization* 47: 35-110.
 - [10] Penot J. P. 1998. "Are generalized derivatives useful for generalized convex functions?", in *Generalized Convexity, Generalized Monotonicity: Recent Results*, J.-P. Crouzeix, J. E. Martinez-Legaz, and M. Volle, (eds.), Kluwer, Dordrecht. 3-59.
 - [11] Penot J. P., Zălinescu C. 2000. "Elements of Quasiconvex Subdifferential Calculus", *Journal of Convex Analysis* 7: 243-269.
 - [12] Rockafellar R. T. 1970. "Convex Analysis", Princeton University Press, Princeton, NJ.

زیر مشتق توپولوژیکی و نقش آن در بهینه‌سازی ناهموار با داده‌های شبه محدب

سروش، ح.

گروه ریاضی، دانشگاه پیام نور، صندوق پستی ۳۶۹۷-۱۹۳۹۵ تهران، ایران،

h_sorosh2011@pnu.ac.ir

تاریخ دریافت: ۲۷ خرداد ۱۴۰۰ تاریخ پذیرش: ۲۸ مهر ۱۴۰۰

چکیده

در این مقاله، با استفاده از زیرمجموعه توپولوژیکی، مسئله‌ی بهینه‌سازی ناهموار را با توابع شبه محدب تجزیه و تحلیل می‌کنیم. در ادامه، برخی از شرایط بهینه‌سازی لازم و کافی را ارائه می‌دهیم و توابع شبه محدب توپولوژی را توصیف می‌کنیم. در انتها، نتایج دوگانگی ضعیف و قوی از نوع موند-ویر را برای این مسئله بیان می‌کنیم.

کلمات کلیدی

بهینه‌سازی شبه محدب، قضیه‌های دوگانی، شرایط بهینگی، شبه محدب، زیرمشتق.